

WEAK SEMISTABLE REDUCTION IN CHARACTERISTIC 0 PRELIMINARY VERSION

DAN ABRAMOVICH[†] AND KALLE KARU

0. INTRODUCTION

Regretfully, we work over an algebraically closed field k of characteristic 0.

0.1. The problem. Roughly speaking, the semistable reduction problem we address here asks for the following:

Let $X \rightarrow B$ be a surjective morphism of complex projective varieties with geometrically integral generic fiber. Find a generically finite proper surjective morphism (that is, an alteration) $B_1 \rightarrow B$, and a proper birational morphism (that is, a modification) $Y \rightarrow X \times_B B_1$, such that the morphism $Y \rightarrow B_1$ is nice.

Of course, one needs to decide what a “nice morphism” means.

The question was posed, among other places, in the introduction of [KKMS], p. vii. It can be viewed as a natural extension of Hironaka’s theorem on resolution of singularities, which is in a sense “the general fiber” of semistable reduction.

0.2. Brief history. The case $\dim B = 1, \dim X = 2$ is very old, see [A-W]. When $\dim B = 1$, semistable reduction was obtained in [KKMS], in the best possible sense: Y is nonsingular, and all the fibers are reduced, strict divisors of normal crossings.

Using a result of Kawamata on ramified covers (see [Kawa], theorem 17), one can obtain semistable reduction “in codimension 1” over a base of arbitrary dimension. Below, we will refer to the result of Kawamata as “**Kawamata’s trick**”. We will discuss it in detail in section 5.

The case where $\dim X = \dim B + 1$ has recently been proven by de Jong [dJ]. Here one shows that any family of curves can be made into a family of nodal curves, which are indeed as “nice” as one may expect.

Using recent difficult results of Alexeev, Kollár and Shepherd-Barron (see [Al], [Al1]), one obtains a version of the case $\dim X = \dim B + 2$. Here each fiber is a semi-log-canonical surface.

Up until recently, not much has been known about the case $\dim X > \dim B + 2$. Often one finds remarks of the following flavor: “since we do not have a semistable reduction result over a base of higher dimension, we will work around it in the following technical manner...”.

0.3. Definition of semistable families. We give here a description of the best possible kind of morphisms we have in mind.

Let $f : X \rightarrow B$ be a flat morphism of nonsingular projective varieties with connected fibers. Somewhat informally, we say that f is **semistable** if for each point $x \in X$ with $f(x) = b$ there is a choice of formal coordinates $\hat{B}_b = \text{Spec } k[[t_i]]$ and $\hat{X}_x = \text{Spec } k[[x_j]]$, such that f is given by:

$$t_i = \prod_{j=l_{i-1}+1}^{l_i} x_j.$$

[†] Partially supported by NSF grant DMS-9700520 and by an Alfred P. Sloan research fellowship.

Here $0 = l_0 < l_1 \cdots < l_m \leq n$, where $n = \dim X$ and $m = \dim B$. To be more precise, we give things a more global structure using the notion of a toroidal morphism. At the same time we describe a slightly weaker condition which will appear below:

Definition 0.1. The morphism $f : X \rightarrow B$ above is called **weakly semistable** if

1. the varieties X and B admit toroidal structures $U_X \subset X$ and $U_B \subset B$, with $U_X = f^{-1}U_B$;
2. with this structure, the morphism f is toroidal;
3. the morphism f is equidimensional;
4. all the fibers of the morphism f are reduced; and
5. B is nonsingular.

If also X is nonsingular, we say that the morphism $f : X \rightarrow B$ is **semistable**.

0.4. The ultimate goal. The result one would really like to have is:

Conjecture 0.2. Let $X \rightarrow B$ be a surjective morphism of complex projective varieties with geometrically integral generic fiber. There is a projective alteration $B_1 \rightarrow B$, and a projective modification $Y \rightarrow X \times_B B_1$, such that $Y \rightarrow B_1$ is semistable.

Naïvely one might hope to have each fiber isomorphic to a divisor of normal crossings. But already in the case of a 2-parameter family of surfaces $t_1 = x_1x_2$; $t_2 = x_3x_4$, this is impossible. It seems that the definition above is the best one can hope for.

0.5. A. J. de Jong's results. In [dJ], Johan de Jong shows, among many other results, that if one allows $Y \rightarrow X \times_B B_1$ to be an alteration instead of a modification, one can make $Y \rightarrow B_1$ very nice indeed: Y is nonsingular, $Y \rightarrow B_1$ is semistable as in the definition above, and moreover it can be written as a composition of nodal curve fibrations $Y = Y_0 \rightarrow Y_1 \rightarrow \cdots \rightarrow Y_k = B_1$.

De Jong's methods and ideas will serve as a starting point for investigating the semistable reduction conjecture.

0.6. Our main result. The main result of this paper is the following:

Theorem 0.3 (Weak semistable reduction). *Let $X \rightarrow B$ be a surjective morphism of complex projective varieties with geometrically integral generic fiber. There exist an alteration $B_1 \rightarrow B$ and a modification $Y \rightarrow X \times_B B_1$, such that $Y \rightarrow B_1$ is weakly semistable.*

With a little more work we will get X to have only quotient singularities. There are many cases (such as when f is a family of surfaces) where we can actually prove the semistable reduction conjecture. These will be pursued elsewhere. Hopefully, by the time this paper achieves its final form the conjecture will be fully proven.

0.7. Mild morphisms. A few words are in order about the significance of our result. Note that the property of a morphism being semistable is far from being stable under base changes. One may ask, what remains from semistability after at least *dominant* base changes? Here is a suggestion:

Definition 0.4. We define a morphism $X \rightarrow B$ as above to be **mild**, if for any dominant $B_1 \rightarrow B$ where B_1 has at most rational Gorenstein singularities, we have that $X \times_B B_1$ has at most rational Gorenstein singularities as well.

Mild morphisms arise naturally in moduli theory. Indeed, mild families of curves are precisely nodal families; families of Gorenstein semi-log-canonical surfaces mentioned above are mild. For

a discussion of why mild morphisms are useful, see [N]. In fact, the paper [N] would have been much simplified, had mild reduction been available.

Already in the case $\dim B = 1$, mild reduction is a much easier task than semistable reduction. Indeed, lemma 2 on page 103 of [KKMS], and the discussion there, already give mild reduction in this case. The delicate combinatorics of chapter III of [KKMS] is not used for this purpose.

It will be shown (see section 6) that weakly semistable morphisms are indeed mild.

0.8. Structure of the proof. After the introduction, section 1 will be devoted to a general discussion of toroidal morphisms. The proof itself will begin with section 2.

Semistable reduction has at least two flavors: first, the fibers of the morphism $Y \rightarrow B_1$ should have nice local defining equations. Second, the family should have nice algebraic properties. We will perform a number of reduction steps, incrementally improving one or the other of these flavors.

0.8.1. Toroidal reduction. In the first step, carried out in section 2, we will show that any morphism can be modified to a toroidal morphism. The construction is inspired by the inductive procedure of [dJ], and follows closely the proofs in [N-dJ].

Just as in [N-dJ], the construction we give is very non-canonical. Even when the generic fiber of $X \rightarrow B$ is smooth, it will be blown up during the construction. One hopes that methods such as those of [B-M] or [Vil] could be adapted to this situation and give a more canonical procedure.

It is tempting to state the following conjecture.

Conjecture 0.5. Let $X \rightarrow B$ be a morphism as in the theorem. Let $U \subset B$ be an open set over which X is toroidal, and let $\Sigma = B \setminus U$. There exists modifications $X' \rightarrow X$ and $B' \rightarrow B$, each of which is the composition of a sequence of blowings up with smooth centers lying over Σ , and a lifting $X' \rightarrow B'$ which is toroidal.

It should be noted, that in view of recent results of Morelli [Mor] and Włodarczyk [Włó], this conjecture implies the strong blow-up - blow-down conjecture.

0.8.2. Improving the toroidal morphism. In sections 3 and 4 we perform a couple of simple reduction steps to improve our situation. Let $f : (U_X \subset X) \rightarrow (U_B \subset B)$ be any toroidal morphism, with B nonsingular. By the results of [KKMS], we can find a toroidal resolution of singularities $X' \rightarrow X$. Let $f' : X' \rightarrow B$ be the resulting projection. We first show that now $f'^{-1}U_B \subset X'$ is also a toroidal embedding, which is easier to handle: there are no horizontal divisors. For convenience, we replace $X \rightarrow B$ by the new morphism. We remark that one can proceed a fair distance without removing these horizontal divisors, and, we believe, the results one can obtain are of interest (e.g., the inductive structure of de Jong can be preserved), but this would make the present paper much more cumbersome, so we delay that investigation to a future occasion.

Now, our morphism $X \rightarrow B$ is not necessarily equidimensional. We repair this by an appropriate decomposition of the associated conical polyhedral complexes Δ_X and Δ_B . We make sure that, after the modification, the base remains nonsingular, and then the morphism is automatically flat.

0.8.3. Kawamata's trick and reduced fibers. We start section 5 with a discussion of Kawamata's trick and its relation with toroidal morphisms in some detail. Then we use Kawamata's trick to find a finite base change, after which all the fibers are reduced. This finishes the proof of the main theorem, since the resulting morphism is weakly semistable. A variant of Kawamata's trick for global "index 1 covers" is discussed in section 7.

0.8.4. *Mild reduction.* We begin section 6 by checking that the resulting fibers are Gorenstein.

Using a base change and descent argument, and the fact that toroidal singularities are always rational, we then prove that the resulting family is mild.

0.8.5. *Combinatorial restatement.* In section 8 the semistable reduction conjecture is restated purely in combinatorial terms. We end the paper with a discussion of the problems one encounters when trying to go from weak semistable reduction to semistable reduction.

0.9. **Acknowledgments.** We would like to thank O. Gabber, A.J. de Jong, H. King and K. Matsuki for helpful and inspiring discussions.

0.10. **Terminology.** A **modification** is a proper birational morphism of irreducible varieties.

An **alteration** $a : B_1 \rightarrow B$ is a proper, surjective, generically finite morphism of irreducible varieties, see [dJ, 2.20]. The alteration a is a **Galois alteration** if there is a finite group $G \subset \text{Aut}_B(B_1)$ such that the associated morphism $B_1/G \rightarrow B$ is birational, compare [dJ2, 5.3].

1. TOROIDAL MORPHISMS

We have collected in this section some notations and preliminaries about toric varieties, toroidal embeddings, and their morphisms (see [KKMS] for details).

Remark 1.1. Our approach here is based on the formalism of [KKMS]. A different approach, using *logarithmic structures*, was developed by K. Kato, see [Kato], [Kato1]. It is our belief, that the approach via logarithmic structures should eventually prevail - it provides us with a flexible category, in which toroidal embeddings (= logarithmically regular schemes) and toroidal morphisms (= logarithmically smooth(!) morphisms) play a special role. Some of our statements below are rendered almost trivial with Kato's formalism, e.g. Lemmas 1.5 and 5.3.

The reason we decided to stick with the formalism of [KKMS] is, that the theory of logarithmic structures is not yet in stable form (see the many flavors of such structures introduced in Kato's papers), and, more importantly, it has not yet gained widespread acceptance as a basic formalism. It might have turned away some readers (especially those combinatorially inclined) had we used the theory of logarithmic structures throughout.

It is also worth noting, that Kato's notion of a fan, although it has a nice structural morphism, is much less amenable to combinatorial manipulation than the polyhedral complexes of [KKMS].

1.1. **Toric varieties.** Given a lattice $N \cong \mathbb{Z}^n$, its dual $M = \text{Hom}(N, \mathbb{Z})$, a strictly convex rational polyhedral cone $\sigma \subset N_{\mathbb{R}} = N \otimes \mathbb{R}$ with its dual $\sigma^\vee = \{m \in M_{\mathbb{R}} \mid m(u) \geq 0 \text{ for all } u \in \sigma\}$, we define the **affine toric variety** $X_\sigma = \text{Spec } S[\sigma]$ where $S[\sigma]$ is the semigroup algebra of $\sigma^\vee \cap M$ over the ground field. If more than one toric variety is considered, we use a subscript: N_σ, M_σ .

We denote by $\sigma^{(1)}$ the 1-dimensional edges of σ . The indivisible points v in $\sigma^{(1)} \cap N$ are called the **primitive points** of σ . The variety X_σ is nonsingular if and only if the primitive points of σ form a part of a basis of N . In that case we say that σ is nonsingular.

The toric variety X_σ contains an n -dimensional algebraic torus $T = \mathbb{G}_m^n$ as an open dense subset, and the action of T on itself extends to an action on X_σ . Thus, X_σ is a disjoint union of orbits of this action. There is a one-to-one correspondence between the orbits and the faces of σ . In particular, 1-dimensional faces $\mathbb{R}_+ v_i$ correspond to codimension 1 orbits $\mathbb{O}_{v_i}$.

A **toric morphism** $f : X_\sigma \rightarrow X_\tau$ is a dominant equivariant morphism of toric varieties defined by a linear map $f_\Delta : (N_\sigma, \sigma) \rightarrow (N_\tau, \tau)$. We use the same notation for the scalar extension $f_\Delta : N_\sigma \otimes \mathbb{R} \rightarrow N_\tau \otimes \mathbb{R}$.

1.2. Toroidal embeddings. Given a normal variety X and an open subset $U_X \subset X$, the embedding $U_X \subset X$ is called **toroidal** if for every closed point $x \in X$ there exist a toric variety X_σ , a point $s \in X_\sigma$, and an isomorphism of complete local algebras

$$\hat{\mathcal{O}}_{X,x} \cong \hat{\mathcal{O}}_{X_\sigma,s}$$

so that the ideal of $X \setminus U_X$ corresponds to the ideal of $X_\sigma \setminus T$. Such a pair (X_σ, s) is called a local model at $x \in X$. By restricting X_σ if necessary, we can assume that the orbit of s is the unique closed orbit in X_σ .

Definition 1.2. A dominant morphism $f : (U_X \subset X) \rightarrow (U_B \subset B)$ of toroidal embeddings is called **toroidal** if for every closed point $x \in X$ there exist local models (X_σ, s) at x , (X_τ, t) at $f(x)$ and a toric morphism $g : X_\sigma \rightarrow X_\tau$ so that the following diagram commutes

$$\begin{array}{ccc} \hat{\mathcal{O}}_{X,x} & \xleftarrow{\cong} & \hat{\mathcal{O}}_{X_\sigma,s} \\ \hat{f}^* \uparrow & & \uparrow \hat{g}^* \\ \hat{\mathcal{O}}_{B,f(x)} & \xleftarrow{\cong} & \hat{\mathcal{O}}_{X_\tau,t} \end{array}$$

where $\hat{f}^*$ and $\hat{g}^*$ are the ring homomorphisms induced by f and g .

1.3. Cones and polyhedral complexes. Let $X \setminus U_X = \cup_{i \in I} E_i$ where E_i are irreducible and have codimension 1. We will assume that all the E_i are normal, that is, $U_X \subset X$ is a **toroidal embedding without self-intersection** (also known as a **strict** toroidal embedding). In that case, we can use the irreducible components of $\cap_{i \in J} E_i$ for all $J \subset I$ to define a stratification of X (these components are the closures of strata). Closures of strata formally corresponds to closures of orbits in local models. Since a toric morphism maps orbits to orbits, a toroidal morphism maps strata to strata.

Let Y be a stratum in X , which is by definition an open set in an irreducible component of $\cap_{i \in J} E_i$ for some $J \subset I$. The star of Y is the union of strata in whose closure Y lies (each of these corresponds to some $K \subset J \subset I$). To the stratum Y we associate

1. M^Y : – the group of Cartier divisors in $\text{Star}(Y)$ supported in $\text{Star}(Y) \setminus U_X$
2. N^Y : – $\text{Hom}(M^Y, \mathbb{Z})$
3. $M_+^Y \subset M^Y$: – effective Cartier divisors
4. $\sigma^Y \subset N_{\mathbb{R}}^Y$: – the dual of M_+^Y

It is shown in [KKMS] (Corollary 1, page 61) that if (X_σ, s) is a local model at $x \in X$ in the stratum Y , then

$$\begin{aligned} M^Y &\cong M_\sigma / \sigma^\perp \\ \sigma^Y &\cong \sigma \end{aligned}$$

The cones σ^Y glue together to form a polyhedral complex $\Delta_X = (|\Delta_X|, \{\sigma^Y\}, \{M^Y\})$, where the lattices M^Y form an integral structure on Δ_X . Equivalently, instead of M^Y we may give the lattices N^Y and embeddings $\sigma^Y \hookrightarrow N_{\mathbb{R}}^Y$. We also denote $M_{\sigma^Y} = M^Y$, $N_{\sigma^Y} = N^Y$. Then, comparing to the lattices of local models,

$$\begin{aligned} M_{\sigma^Y} &\cong M_\sigma / \sigma^\perp \\ N_{\sigma^Y} &\cong N_\sigma \cap \text{Span}(\sigma) \end{aligned}$$

If $N_{\sigma^Y} \neq N_\sigma$ we can choose a splitting of N_σ so that at a point $x \in Y$ the local model is in the form $(X_{\sigma'} \times \mathbb{G}_m^l, (s', 1))$ where $(N_{\sigma'}, \sigma') \cong (N_{\sigma^Y}, \sigma^Y)$.

Lemma 1.3. *A toroidal morphism $f : X \rightarrow B$ induces a morphism $f_\Delta : \Delta_X \rightarrow \Delta_B$, which for each cone σ^Y is the restriction of $g_\Delta : (\sigma, N_\sigma) \rightarrow (\tau, N_\tau)$ where (X_σ, s) , (X_τ, t) are local models at $x \in Y \subset X$, $f(x) \in Z \subset B$, and g_Δ is the linear map determined by the toric morphism $g : X_\sigma \rightarrow X_\tau$ in Definition 1.2.*

Proof. To see that the maps f_Δ defined for different cones σ^Y agree on the overlaps it suffices to notice that the dual morphism $f_\Delta^\vee : M_{\tau Z} \rightarrow M_{\sigma Y}$ is defined by pulling back a Cartier divisor and restricting it to $\text{Star}(Y) \setminus U_X$. Since the pullback is defined independently of the stratum, we see that f_Δ is well defined. $\square$

Remark 1.4. Note that the polyhedral morphism $f_\Delta : \Delta_X \rightarrow \Delta_B$ is well defined even if f is not toroidal, as long as $f(U_X) \subset U_B$.

Lemma 1.5. *If $e : X \rightarrow Y$ and $f : Y \rightarrow Z$ are toroidal morphisms, then $f \circ e : X \rightarrow Z$ is also toroidal.*

Remark 1.6. This lemma is a triviality if one uses logarithmic structures.

Proof. Let $x \in X$, $y = e(x) \in Y$, $z = f(y) \in Z$, and choose local models at x , y and z as in Definition 1.2. Consider the tower

$$\begin{array}{ccc}
 \hat{\mathcal{O}}_{X,x} & \xleftarrow{\cong} & \hat{\mathcal{O}}_{X_\sigma,s} \\
 \hat{e}^* \uparrow & & \uparrow \hat{g}^* \\
 \hat{\mathcal{O}}_{Y,y} & \xleftarrow{\cong} & \hat{\mathcal{O}}_{X_{\tau_1},t_1} \\
 id \uparrow & & \uparrow \alpha \\
 \hat{\mathcal{O}}_{Y,y} & \xleftarrow{\cong} & \hat{\mathcal{O}}_{X_{\tau_2},t_2} \\
 \hat{f}^* \uparrow & & \uparrow \hat{h}^* \\
 \hat{\mathcal{O}}_{Z,z} & \xleftarrow{\cong} & \hat{\mathcal{O}}_{X_\rho,r}
 \end{array}$$

where the upper and lower squares commute by the definition of toroidal morphism, and where α is defined by tracing the other three sides of the middle square. Then the middle square also commutes.

Since τ_1 and τ_2 are isomorphic, we can take $\tau_1 = \tau_2 = \tau$ and $X_{\tau_1} = X_{\tau_2} = X_\tau$. The map α , of course, need not be the identity. Let the coordinate rings of the tori in X_σ , X_τ and X_ρ be $k[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}]$, $k[y_1, y_1^{-1}, \dots, y_m, y_m^{-1}]$ and $k[z_1, z_1^{-1}, \dots, z_l, z_l^{-1}]$, respectively, so that the toric morphisms g and h are defined by

$$y_i \mapsto \prod_{j=1}^n x_j^{a_{ij}}, \quad z_i \mapsto \prod_{j=1}^m y_j^{b_{ij}}$$

for some $a_{ij}, b_{ij} \in \mathbb{Z}$.

The maps in the third line of the tower identify the group of T-invariant Cartier divisors in X_{τ_2} with the group of Cartier divisors in Y supported in $Y \setminus U_Y$ and passing through y . The maps in the second line of the tower identify the latter group with the group of T-invariant Cartier divisors in X_{τ_1} . Hence α induces a group homomorphism between T-invariant Cartier divisors of X_{τ_2} and X_{τ_1} . Since T-invariant Cartier divisors in X_τ are given by products of y_i ,

then (using the same letter y_i for the image of y_i in the completed local ring) α maps

$$y_i \mapsto u_i \prod_{j=1}^m y_j^{c_{ij}}$$

where u_i are units in $\hat{\mathcal{O}}_{X_\tau, t_1}$, and $c_{ij} \in \mathbb{Z}$.

The composition $\hat{g}^* \circ \alpha \circ \hat{h}^*$ is then defined by

$$z_i \mapsto v_i \prod_{j=1}^m x_i^{d_{ij}}$$

where the matrix with entries d_{ij} is the product of the matrices with entries a_{ij} , c_{ij} and b_{ij} , and where $v_i = \hat{g}^*(\prod_j u^{c_{ij}})$ are units in $\hat{\mathcal{O}}_{X_\sigma, s}$. The matrix (d_{ij}) is equivalent to a matrix in the form

$$\begin{pmatrix} D \\ 0 \end{pmatrix}$$

where D is an $l \times l$ diagonal matrix with diagonal entries $d_1, \dots, d_l \in \mathbb{Z}$. Hence we can change the coordinate functions x_i and z_i of the respective tori so that $\hat{g}^* \circ \alpha \circ \hat{h}^*$ maps

$$z_i \mapsto v_i x_i^{d_i}$$

If we now set

$$\tilde{x}_i = v_i^{\frac{1}{d_i}} x_i$$

then in these new coordinates the composition $\hat{g}^* \circ \alpha \circ \hat{h}^*$ is induced by a toric morphism defined by

$$z_i \mapsto x_i^{d_i}$$

and thus $f \circ e : X \rightarrow Z$ is toroidal. $\square$

Given a toroidal embedding $U_X \subset X$ with polyhedral complex Δ_X , and a subdivision Δ'_X of Δ_X , one constructs (see [KKMS]) a new toroidal embedding $U_{X'} \subset X'$ with polyhedral complex Δ'_X , and a birational toroidal morphism $f' : X' \rightarrow X$ such that the induced map of the polyhedral complexes $\Delta'_X \rightarrow \Delta_X$ is the given subdivision. If Y is a stratum in X corresponding to the cone $\sigma^Y \in \Delta_X$, and if $\sigma' \subset \sigma$ is a cone in the subdivision, define

$$V_{\sigma'} = \text{Spec } \text{Star}(Y) \sum_{D \in \sigma'^{\vee} \cap M^Y} \mathcal{O}_{\text{Star}(Y)}(-D),$$

where the sum is taken inside the field of rational functions of $\text{Star}(Y)$. Then X' is formed by gluing together the open sets $V_{\sigma'}$.

A subdivision Δ'_X of Δ_X is called **projective** if there exists a continuous function $\psi : |\Delta_X| \rightarrow \mathbb{R}$, taking rational values on $\sigma \cap N_\sigma$, which is convex and piecewise linear on each cone $\sigma \in \Delta_X$, and the largest pieces in σ where ψ is linear are the cones of the subdivision. Such ψ is called a **good function** (or *lifting function*, or *order function*), for the subdivision Δ'_X of Δ_X . A projective subdivision corresponds to a projective modification $X' \rightarrow X$.

Lemma 1.7. *Let $f : X \rightarrow B$ be a toroidal morphism, $f_\Delta : \Delta_X \rightarrow \Delta_B$ the associated morphism of polyhedral complexes. Let $X' \rightarrow X$ and $B' \rightarrow B$ be toroidal modifications, with associated subdivisions $\Delta_{X'}$ and $\Delta_{B'}$. Then f lifts to a morphism $f' : X' \rightarrow B'$ if and only if for each cone $\sigma' \in \Delta_{X'}$, there exists a cone $\tau' \in \Delta_{B'}$ such that $f_\Delta(\sigma') \subset \tau'$.*

Proof. Let $\sigma' \subset \sigma^Y \in \Delta_X$ and $\tau' \subset \tau^Z \in \Delta_B$ be cones in the subdivisions such that $f_\Delta(\sigma') \subset \tau'$. The homomorphism $\mathcal{O}_{\text{Star}(Z)} \rightarrow \mathcal{O}_{\text{Star}(Y)}$ extends to

$$\sum_{D \in \tau'^\vee \cap M^Z} \mathcal{O}_{\text{Star}(Z)}(-D) \rightarrow \sum_{E \in \sigma'^\vee \cap M^Y} \mathcal{O}_{\text{Star}(Y)}(-E)$$

because $f_\Delta^\vee(\tau'^\vee) \subset \sigma'^\vee$. This shows that the rational map f' is a morphism on each $V_{\sigma'}$. Since $V_{\sigma'}$ cover X' , f' is a morphism.

Conversely, suppose f' is a morphism. Let $x' \in X'$, $f'(x') = b' \in B'$. Since the three morphisms $X' \rightarrow X$, $X \rightarrow B$ and $B' \rightarrow B$ are toroidal, we have a toric morphism $g : X_{\sigma'} \rightarrow X_{\tau'}$ where $(X_{\sigma'}, s')$ and $(X_{\tau'}, t')$ are local models at x' and b' . Thus $g_\Delta(\sigma') \subset \tau'$. But $g_\Delta|_{\sigma'} = f_\Delta|_{\sigma'}$, hence $f_\Delta(\sigma') \subset \tau'$. Since this is true for x' in any stratum of X , we get that for any cone $\sigma' \in \Delta_{X'}$, $f_\Delta(\sigma') \subset \tau'$ for some $\tau' \in \Delta_{B'}$. $\square$

2. TOROIDAL REDUCTION

2.1. Statement of result. The purpose of this section is to modify any family of varieties into a toroidal morphism.

Theorem 2.1. *Let $f : X \rightarrow B$ be a projective, surjective morphism with geometrically integral generic fiber, and assume B integral. Let $Z \subset X$ be a proper closed subscheme. There exist a diagram as follows:*

$$\begin{array}{ccccc} U_X & \subset & X' & \xrightarrow{m_X} & X \\ \downarrow & & \downarrow f' & & \downarrow f \\ U_B & \subset & B' & \xrightarrow{m_B} & B \end{array}$$

such that m_B and m_X are modifications, X' and B' are nonsingular, the inclusions on the left are toroidal embeddings, and such that

1. f' is toroidal.
2. Let $Z' = m_X^{-1}Z$. Then Z' is a strict normal crossings divisor, and $Z' \subset X' \setminus U_{X'}$.

2.2. To begin the proof, we proceed by induction on the relative dimension of f .

If the relative dimension of f is 0, let $m_X : X' \rightarrow X$ be a resolution of singularities such that $Z' = m_X^{-1}Z$ is a strict normal crossings divisor, let $B' = X'$ and $m_B = f \circ m_X$, and $f' = id$ the identity.

Assume we have proven the result for morphisms of relative dimension $n - 1$, and consider the case $\text{rel. dim } f = n$.

2.3. Preliminary reduction steps. First, we may replace X by its normalization, therefore we may assume X normal, and by blowing up Z in X we may assume Z a Cartier divisor. Let $\eta \in B$ be the generic point of B . By the projectivity assumption we have $X \subset \mathbb{P}_B^N$ for some N . Choosing a generic projection $\mathbb{P}_\eta^N \dashrightarrow \mathbb{P}_\eta^{n-1}$ we get a rational map $X_\eta \dashrightarrow \mathbb{P}_\eta^{n-1}$. Replacing X by the closure of the graph of this map, we may assume that we have a morphism $g : X \rightarrow \mathbb{P}_B^{n-1} = P$.

2.4. Semistable reduction of a family of curves. By [dJ2], Theorem 2.4, we have a diagram as follows:

$$\begin{array}{ccccc}
X_1 & \xrightarrow{\alpha} & X & & \\
\downarrow g_1 & & \downarrow g & & \\
P_1 & \xrightarrow{a} & P & & \\
& & \downarrow & & \\
& & B & &
\end{array}$$

and a finite group $G \subset \text{Aut}_P P_1$, with the following properties:

1. The morphism $a : P_1 \rightarrow P$ is a Galois alteration with Galois group G .
2. The action of G lifts to $\text{Aut}_X X_1$, and $\alpha : X_1 \rightarrow X$ is a Galois alteration with Galois group G .
3. There are n disjoint sections $\sigma_i : P_1 \rightarrow X_1$ such that the strict altered transform $Z_1 \subset X_1$ of Z is the union of their images, and G permutes the sections σ_i .
4. The morphism $g_1 : X_1 \rightarrow P_1$ is a nodal family of curves, and $\sigma_i(P_1)$ is disjoint from $\text{Sing } g_1$.

We may replace X , P and Z by X_1/G and P_1/G , and $\alpha^{-1}Z/G$. Note that $\alpha^{-1}Z/G$ is not necessarily equal to the union of the images of σ_i , but the complement lies over a proper closed subset in P_1 .

2.5. Using the inductive hypothesis. Let $\Delta \subset P$ be the union of the loci over which Z, P_1 or X_1 are not smooth. We apply the inductive assumption to $\Delta \subset P \rightarrow B$, and obtain a diagram as follows:

$$\begin{array}{ccccc}
U_P & \hookrightarrow & P' & \xrightarrow{m} & P \\
\downarrow & & \downarrow & & \downarrow \\
U_B & \hookrightarrow & B' & \rightarrow & B
\end{array}$$

Such that P', B' are nonsingular, $P' \rightarrow P$ and $B' \rightarrow B$ are modifications, the left square is a toroidal morphism, and $m^{-1}\Delta$ is a divisor of strict normal crossings contained in $P' \setminus U_P$.

We may again replace P, B by P', B' , and further we may replace X, X_1, P_1, Z and σ_i by their pullback to P' . In particular $P \rightarrow B$ has a toroidal structure, and $P_1 \rightarrow P$ is unramified over U_P . By Abhyankar's lemma, since P_1 is normal, it inherits a toroidal structure given by $U_{P_1} = m^{-1}U_P$ as well, so that $P_1 \rightarrow P$ is a toroidal finite morphism.

2.6. Conclusion of proof. Now $X_1 \rightarrow P_1$ is a nodal family which is smooth over U_P , therefore it as well inherits a toroidal structure $U_{X_1} \subset U_X$, where $U_{X_1} = (g_1^{-1}U_{P_1}) \setminus (\cup \sigma_i(P_1))$; e.g. local equations around a node are of the form $uv = f(t)$, where $f(t)$ is a monomial on P_1 . Notice that $\alpha^{-1}Z$ is a divisor contained in U_{X_1} (see [N-dJ], 1.3).

In this situation we can apply the procedure of [N-dJ], section 1.4 to make the group G act toroidally on X_1 : first we blow up the scheme $\text{Sing } g_1$ to separate the branches of the nodes. Then we are in the situation of Proposition 1.8 of [N-dJ], namely there is a canonical G -equivariant blowup $d : \tilde{X}_1 \rightarrow X_1$ such that G acts strictly toroidally on $b^{-1}U_{X_1} \subset \tilde{X}_1$. Let $X' = \tilde{X}_1/G$, then $X' \rightarrow B$ inherits a toroidal structure and $X' \rightarrow X$ is birational; moreover, $Z' \subset X'$ is a divisor contained in $X' \setminus U_{X'}$. Applying toroidal resolution of singularities, the induction step is proven. $\square$

3. REMOVING HORIZONTAL DIVISORS

We may now replace $X \rightarrow B$ by $X' \rightarrow B'$, and thus we may assume that the morphism f is toroidal. Our goal in this section is to arrive at a situation where $f^{-1}U_B = U_X$.

The rough idea is, that a morphism between nonsingular toroidal embeddings $f : X \rightarrow B$ is locally given by monomials $t_i = x_1^{k_1} \cdots x_r^{k_r}$, in which the variables defining horizontal divisors

cannot appear, so these divisors are unnecessary in the toroidal description. We make this precise by a simple translation argument.

Proposition 3.1. *Let $U_X \subset X$ and $U_B \subset B$ be nonsingular toroidal embeddings and $f : X \rightarrow B$ a surjective toroidal morphism. Then, denoting $U'_X = f^{-1}(U_B) \supset U_X$, we have that $U'_X \subset X$ is a toroidal embedding, and $f : (U'_X \subset X) \rightarrow (U_B \subset B)$ is a toroidal morphism.*

Proof. Since f maps U_X into U_B , $f^{-1}(B \setminus U_B)$ as a set is a union of divisors supported in $X \setminus U_X$. In local models these divisors are all T-invariant.

Consider local models (X_σ, s) at x , (X_τ, t) at $f(x)$, and the toric morphism $g : X_\sigma \rightarrow X_\tau$. We may assume that $v_1, \dots, v_n$ is a basis of N_σ and σ is generated by $v_1, \dots, v_k$. Then $X_\sigma \cong \mathbb{A}^k \times \mathbb{G}_m^{n-k}$, and we may take $s = (0, 1)$. Let the closures of the orbits corresponding to $v_1, \dots, v_j$ be the horizontal divisors. That means, $g_\Delta(v_i) = 0$ for $i = 1, \dots, j$, and g factors through the projection:

$$g : X_\sigma \cong \mathbb{A}^j \times \mathbb{A}^{k-j} \times \mathbb{G}_m^{n-k} \rightarrow \mathbb{A}^{k-j} \times \mathbb{G}_m^{n-k} \rightarrow X_\tau$$

Now take $s' = (1, 0, 1) \in \mathbb{A}^j \times \mathbb{A}^{k-j} \times \mathbb{A}^{n-k}$. From the factorization of g we see that $g(s') = g(s)$. Translation by $(1, 0, 0)$ gives an isomorphism of the local rings at s and s' so that the ideals of the T-invariant divisors passing through s' corresponds to the ideal of the vertical T-invariant divisors passing through s . Thus, (X_σ, s') is a local model for the embedding $U'_X \subset X$ at $x \in X$ and $g : (X_\sigma, s') \rightarrow (X_\tau, t)$ is the toric morphism of the local models representing f . $\square$

4. MAKING THE MORPHISM EQUIDIMENSIONAL

The goal of this section is to perform modifications on B and X , after which the morphism becomes equidimensional. First a lemma which characterizes equidimensional toroidal morphisms:

Lemma 4.1. *Let $f : X \rightarrow B$ be a surjective toroidal morphism, $f_\Delta : \Delta_X \rightarrow \Delta_B$ the associated morphism of polyhedral complexes. Then f is equidimensional if and only if for any cone $\sigma \in \Delta_X$, we have $f_\Delta(\sigma) \in \Delta_B$. That is, the image of a cone of Δ_X is a cone of Δ_B .*

Proof. Computing the dimension of a local ring commutes with taking the completion. Thus, it suffices to consider local models. The generic fiber of a toric morphism $f : X_\sigma \rightarrow X_\tau$ has dimension $\text{Rank}(N_\sigma) - \text{Rank}(N_\tau)$. Now f maps a k -dimensional orbit corresponding to a $(\text{Rank}(N_\sigma) - k)$ -dimensional face σ' of σ onto some l -dimensional orbit corresponding to a $(\text{Rank}(N_\tau) - l)$ -dimensional face τ' of τ . Thus f is equidimensional if and only if

$$(\text{Rank}(N_\sigma) - k) - (\text{Rank}(N_\tau) - l) \leq \text{Rank}(N_\sigma) - \text{Rank}(N_\tau),$$

that means $l \leq k$. Every k -dimensional cone maps to an l -dimensional cone for $l \leq k$ if and only if the image of every cone is a cone. $\square$

Remark 4.2. In case τ is simplicial the condition of the theorem is equivalent to the statement that all 1-dimensional faces of σ map to 0 or 1-dimensional faces of τ .

The following lemma is a slight generalization of the toric Chow's lemma ([D] 6.9.2 page 119).

Lemma 4.3. *Given a polyhedral complex Δ and a subdivision Δ' of Δ , there exists a projective subdivision Δ'' of Δ which refines Δ' .*

Proof. First, we show that it suffices to find a “good” function ψ (see Section 1) on each cone $\sigma \in \Delta$. Indeed, if for each $\sigma \in \Delta$ we have found a good function $\psi_\sigma : \sigma \rightarrow \mathbb{R}$, then a good function on $|\Delta|$ is the sum

$$\psi = \sum_{\sigma \in \Delta} \bar{\psi}_\sigma$$

where $\bar{\psi}_\sigma : |\Delta| \rightarrow \mathbb{R}$ is a good extension of ψ_σ constructed as follows. To extend ψ_σ to a cone $\tau \in \Delta$ we proceed by induction on the dimension of τ (cf. [KKMS] Lemma 1, page 33). If $\dim \tau = 1$ and τ is not a face of σ , define $\bar{\psi}_\sigma|_\tau \equiv 0$. Now assume that $\dim \tau > 1$ and $\bar{\psi}_\sigma$ is defined on $\partial\tau$. Choose a point x in the relative interior of τ and define

$$\bar{\psi}_\sigma(\lambda x + (1 - \lambda)y) = \lambda C + (1 - \lambda)\bar{\psi}_\sigma(y), \quad y \in \partial\tau, 0 \leq \lambda \leq 1.$$

For big $C \in \mathbb{Q}$ the extension $\bar{\psi}_\sigma|_\tau$ is convex.

Now let $\sigma \in \Delta$ with $\dim \sigma = n$. For every $n - 1$ dimensional cone $\tau \in \Delta'$ in the subdivision of σ choose a linear rational function $l_\tau : \sigma \rightarrow \mathbb{R}$ such that τ is the subset of σ defined by $l_\tau = 0$. Then the sum

$$\psi_\sigma(x) = - \sum_{\tau} |l_\tau(x)|$$

is a good function on σ . □

Proposition 4.4. *Let $U_X \subset X$ and $U_B \subset B$ be toroidal embeddings with polyhedral complexes Δ_X and Δ_B respectively, and assume that B is nonsingular. Let $f : X \rightarrow B$ be a surjective toroidal morphism. Then there exist projective subdivisions Δ'_X of Δ_X and Δ'_B of Δ_B with Δ'_B nonsingular, such that the induced map $f' : X' \rightarrow B'$ is an equidimensional toroidal morphism. If $f^{-1}(U_B) = U_X$ then $(f')^{-1}(U_{B'}) = U_{X'}$.*

Proof. There exists a subdivision of Δ_B “induced” by Δ_X . For $x \in \tau \in \Delta_B$ let S_x be the set of cones $\sigma \in \Delta_X$ such that $\sigma \cap f_\Delta^{-1}(x) \neq \{0\}$. Since f_Δ is surjective, $S_x \neq \emptyset$. The relation $x \sim y \Leftrightarrow S_x = S_y$ for $x, y \in \tau$ is clearly an equivalence, hence it defines a partition of Δ_B . The equivalence class of x is

$$\bigcap_{\sigma \in S_x} f_\Delta(\sigma)$$

which is a convex rational polyhedral subcone of τ . Thus the partition defines a subdivision Δ_B^0 of Δ_B such that $f_\Delta(\sigma)$ for any cone $\sigma \in \Delta_X$ is a union of cones in Δ_B^0 . By the previous lemma there exists a refinement Δ_B^1 of Δ_B^0 which is a projective subdivision of Δ_B . Finally, we let Δ'_B be a nonsingular projective subdivision of Δ_B^1 .

For f_Δ to map cones of Δ_X to cones of Δ'_B , the complex Δ_X has to be subdivided. Since the subdivision Δ'_B of Δ_B is projective, there exists a good function ψ on $|\Delta'_B| = |\Delta_B|$. The piecewise linear function $\psi \circ f_\Delta$ defines a projective subdivision Δ'_X of Δ_X whose cones map into cones of Δ'_B . If $\sigma \in \Delta'_X$ then $f_\Delta(\sigma)$ is a union of faces of some cone $\tau \in \Delta_B$. Since f_Δ is linear on σ , $f_\Delta(\sigma)$ is convex in τ , hence $f_\Delta(\sigma)$ is a face of τ .

If we assume from the beginning that $f^{-1}(U_B) = U_X$, that means $f_\Delta^{-1}(0) \cap |\Delta_X| = 0$, then clearly the same is true for any subdivision $|\Delta'_X| = |\Delta_X|$, hence $(f')^{-1}(U_{B'}) = U_{X'}$. □

5. KAWAMATA’S TRICK AND REDUCED FIBERS

5.1. Statement of result. The goal in this section is to find a finite base change, after which all the fibers in the resulting morphism are reduced. The base change we perform will not necessarily be toroidal, but the morphism after base change will still be toroidal.

Proposition 5.1. *Let $U_X \subset X$ and $U_B \subset B$ be projective toroidal embeddings, and assume that B is nonsingular. Let $f : X \rightarrow B$ be a surjective equidimensional toroidal morphism with $f^{-1}(U_B) = U_X$. Then there exists a finite surjective morphism $p : B' \rightarrow B$ so that, denoting by X' the normalization of $X \times_B B'$, we have that B' and X' are toroidal embeddings, the projection $f' : X' \rightarrow B'$ is an equidimensional toroidal morphism with reduced fibers, and $(f')^{-1}(U_{B'}) = U_{X'}$.*

The construction of X' and B' is more explicitly given in Proposition 5.9, where the polyhedral complexes of X' and B' are also described.

5.2. The toric pictures. To start, we characterize equidimensional toroidal morphisms with reduced fibers in terms of polyhedral complexes.

Lemma 5.2. *Let $f : X \rightarrow B$ be an equidimensional toroidal morphism, $f_\Delta : \Delta_X \rightarrow \Delta_B$ the associated morphism of polyhedral complexes. Then f has reduced fibers if and only if for any cone $\sigma \in \Delta_X$, with image $\tau \in \Delta_B$, we have $f_\Delta(N_\sigma \cap \sigma) = N_\tau \cap \tau$. That is, the image of the lattice in any cone of X is the lattice in the image cone.*

Proof. It suffices to consider the toric morphism of local models $f : X_\sigma \rightarrow X_\tau$ and the fiber over a point $t \in X_\tau$ lying in the closed orbit of X_τ . If the orbit of t is not $\{t\}$ then X_τ is a product $X_\tau = X_{\tau'} \times \mathbb{G}_m^q$ for some $q > 0$. Without loss of generality we may then replace X_τ by $X_{\tau'}$, and replace f by $p \circ f$ where $p : X_\tau \rightarrow X_{\tau'}$ is the projection. Indeed, the fiber $(p \circ f)^{-1}(p(t))$ is isomorphic to $f^{-1}(t) \times \mathbb{G}_m^q$, and p_Δ gives an isomorphism $p_\Delta : N_\tau \cap \tau \cong N_{\tau'} \cap \tau'$. Thus we may assume that $\{t\}$ is the unique closed orbit of X_τ . The ideal of $f^{-1}(t)$ is generated by $k[f_\Delta^\vee(\tau^\vee \cap (M_\tau \setminus \{0\}))] \subset k[\sigma^\vee \cap M_\sigma]$. The fiber is reduced if and only if the image $f_\Delta^\vee(\tau^\vee \cap (M_\tau \setminus \{0\}))$ is saturated in $\sigma^\vee \cap M_\sigma$. This happens if and only if $f_\Delta(N_\sigma \cap \sigma) = N_\tau \cap \tau$. $\square$

When X_τ is nonsingular, the condition of the lemma is that primitive points of σ map to primitive points of τ .

Lemma 5.3. *Let $X_\sigma \rightarrow X_\tau$ be a toric morphism with X_τ nonsingular. Let $X_{\tau'}$ be a toric variety given by $\tau' = \tau$ and $N_{\tau'} \subset N_\tau$ a sublattice of finite index. Then every irreducible component of the normalization of $X_\sigma \times_{X_\tau} X_{\tau'}$ is a toric variety $X_{\sigma'}$ given by the cone $\sigma' = \sigma$ and integral lattice $N_{\sigma'} = N_\sigma \cap f_\Delta^{-1}(N_{\tau'})$.*

Proof. The ring of regular functions of $X_{\tau'} \times_{X_\tau} X_\sigma$ is

$$\mathcal{O}_{X_{\tau'}} \otimes_{\mathcal{O}_{X_\tau}} \mathcal{O}_{X_\sigma} = k[(\tau')^\vee \cap M_{\tau'}] \otimes_{k[\tau^\vee \cap M_\tau]} k[\sigma^\vee \cap M_\sigma] = k[\pi]$$

where π is the pushout of $j : \tau^\vee \cap M_\tau \rightarrow (\tau')^\vee \cap M_{\tau'}$ and $f_\Delta^\vee : \tau^\vee \cap M_\tau \rightarrow \sigma^\vee \cap M_\sigma$:

$$\pi = ((\tau')^\vee \cap M_{\tau'}) \times (\sigma^\vee \cap M_\sigma) / \sim.$$

Here $\sim$ is the equivalence relation generated by:

$(v_1, w_1) \sim (v_2, w_2)$ whenever there exists $u \in \tau^\vee \cap M_\tau$ such that $(v_1, w_1) = (v_2, w_2) \pm (u, -f_\Delta^\vee(u))$.

Let M be the abelian group

$$M = M_{\tau'} \times M_\sigma / ((u, -f_\Delta^\vee(u)) | u \in \tau^\vee \cap M_\tau).$$

We will show that the semigroup homomorphism $\iota : \pi \rightarrow M$ is injective. Suppose that $\iota(v_1, w_1) = \iota(v_2, w_2)$, where $v_1, v_2 \in (\tau')^\vee \cap M_{\tau'}$ and $w_1, w_2 \in \sigma^\vee \cap M_\sigma$. Say $(v_1, w_1) = (v_2, w_2) + (u, -f_\Delta^\vee(u))$ for some $u \in M_\tau$. Let $t_1, \dots, t_m \in \tau^\vee$ be a basis of M_τ , so that in this basis $\tau^\vee$ is given by the inequalities $t_i \geq 0$, for $i = 1, \dots, k$. Write $u = \sum_i \alpha_i t_i$ with $\alpha_i \in \mathbb{Z}$. Collecting positive and negative terms, we get $u = u_1 - u_2$ with $u_1, u_2 \in \tau^\vee \cap M_\tau$. Writing v_1, v_2

in terms of t_i and expanding the equation $v_1 = v_2 + u_1 - u_2$, we see that $v_2 - u_2 \in \tau^\vee \cap M_{\tau'}$. Let $(v_3, w_3) = (v_2 - u_2, w_2 + f_\Delta^\vee(u_2))$. Clearly

$$(v_1, w_1) \sim (v_3, w_3) \sim (v_2, w_2)$$

and ι is injective.

The integral closure of $k[\pi]$ in $k[M]$ is the semigroup algebra $k[\tilde{\pi}]$ where $\tilde{\pi}$ is the saturation of π in M . Write $M = F \oplus T$ where F is the free part and T is torsion. Then $T \subset \tilde{\pi}$ because $mT = 0$ for some $m > 0$. For any $f + t \in \tilde{\pi}$ where $f \in F$ and $t \in T$, we have $-t \in T \subset \tilde{\pi}$, hence $f \in \tilde{\pi}$. Thus

$$\tilde{\pi} \cong (\tilde{\pi} \cap F) \oplus T.$$

If t_i are generators of T of order m_i , and if x_i is the image of t_i in $k[\tilde{\pi}]$, then

$$k[\tilde{\pi}] \cong k[\tilde{\pi} \cap F][\dots, x_i, \dots] / (x_i^{m_i} = 1)$$

Thus the normalization of $\text{Spec } k[\tilde{\pi}]$ has $|T|$ components with $x_i = \zeta_i$ where ζ_i are m_i 'th roots of unity, each component isomorphic to $\text{Spec } k[\tilde{\pi} \cap F]$.

Next we show that F can be embedded in $M_\sigma \otimes \mathbb{R}$ so that $M_\sigma \subset F$. The image of the homomorphism $\phi : M \rightarrow M_\sigma \otimes \mathbb{R}$ defined by $\phi(v, w) = f_\Delta^\vee(v) + w$ for $v \in M_{\tau'}, w \in M_\sigma$ contains M_σ . Since M and M_σ have the same rank, ϕ embeds F in $M_\sigma \otimes \mathbb{R}$ as a lattice of full rank containing M_σ . As $f_\Delta^\vee$ takes $\tau^\vee$ into $\sigma^\vee$, we see that ϕ maps $\tilde{\pi} \cap F$ into $\sigma^\vee$ so that the image contains $\sigma^\vee \cap M_\sigma$. Therefore, $\text{Spec } k[\tilde{\pi} \cap F]$ is a toric variety defined by the cone $\sigma' = \sigma$ and integral lattice $N_{\sigma'} = F^\vee$. To determine $F^\vee$, first, we have $M_\sigma \subset F$, hence $F^\vee \subset N_\sigma$; second, $f_\Delta^\vee(M_{\tau'}) \subset F$ implies that $f_\Delta(F^\vee) \subset N_{\tau'}$. Conversely, $N_\sigma \cap f_\Delta^{-1}(N_{\tau'}) \subset F^\vee$, so the two are equal. $\square$

To get a toric morphism with reduced fibers, one can take a base change $X_{\tau'} \rightarrow X_\tau$, where $\tau' = \tau$ and $N_{\tau'} \subset N_\tau$ is a sublattice of finite index. By Lemma 5.3 every component $X_{\sigma'}$ of the normalization of $X_\sigma \times_{X_\tau} X_{\tau'}$ is a toric variety defined by the cone $\sigma' = \sigma$ and integral lattice $N_{\sigma'} = N_\sigma \cap f_\Delta^{-1}(N_{\tau'})$. By a judicious choice of $N_{\tau'}$ the fibers of $X_{\sigma'} \rightarrow X_{\tau'}$ are reduced.

5.3. Kawamata's covering. To perform a similar base change in the toroidal case we need a toroidal morphism $B' \rightarrow B$ which ramifies over a divisor D with a certain index.

Definition 5.4. Let L be a Cartier divisor on B , and let $D \in |mL|$. Choose a rational section s_L of $\mathcal{O}_B(L)$ defining L and a regular section s_D of $\mathcal{O}_B(mL)$ defining D . Consider the rational function $\phi = s_D/s_L^m$ on B . Then the field $K(B)(\sqrt[m]{\phi})$ depends only on D and $\mathcal{O}(L)$. The normalization of B in $K(B)(\sqrt[m]{\phi})$ is called **the cyclic cover ramified along D with index m** .

Remark 5.5. Another way to define the cyclic cover, is as the normalization of

$$\text{Spec}_B \text{Sym}^\bullet(\mathcal{O}_B(L)^\vee) / (\{f - s(f)\}_f)$$

where we view s as a morphism $s : \mathcal{O}_B(kL) \rightarrow \mathcal{O}_B((k+m)L)$.

When B is nonsingular and D a divisor of normal crossings so that $s_D = x_1 \cdots x_l$ for some local parameters $x_1, \dots, x_l$, then the cyclic cover has a local equation

$$z^m = x_1 \cdots x_l.$$

It is nonsingular if and only if $l = 1$.

Let $U_B \subset B$ be a nonsingular projective (strict) toroidal embedding. Then $B \setminus U_B = \sum D_i$ is a strict divisor of normal crossings. Consider the data (D_i, m_i) where D_i are the irreducible components of $B \setminus U_B$ and m_i are positive integers, $i = 1, \dots, m$.

A *Kawamata covering package* consists of (D_i, m_i, H_{ij}) with the following properties:

1. H_{ij} are effective reduced nonsingular divisors on B , for $i = 1, \dots, m$, $j = 1, \dots, \dim B$.
2. $\sum_i D_i + \sum_{i,j} H_{ij}$ is a reduced divisor of normal crossings (in particular, H_{ij} are distinct).
3. $D_i + H_{ij} \in |m_i L_i|$ for some Cartier divisor L_i for all i, j .

To find H_{ij} , let M be an ample divisor. Take a multiple of M if necessary so that $m_i M - D_i$ is very ample for all i , and choose H_{ij} general members in $|m_i M - D_i|$.

Now let B_{ij} be the cyclic cover ramified along $D_i + H_{ij}$ with index m_i ; let B' be the normalization of

$$B_{1,1} \times_B \dots \times_B B_{m,\dim B}$$

and let $p : B' \rightarrow B$ be the projection.

Lemma 5.6 (Kawamata). *The variety B' is nonsingular, ramified with index m_i along $D_i + H_{ij}$. The reduction of the inverse image $p^*(\sum_i D_i + \sum_{i,j} H_{ij})_{\text{red}}$ is a divisor of normal crossings.*

Proof. Let x_i be local equations of D_i , and let y_{ij} be local equations of H_{ij} at $b \in B$. Then B_{ij} is locally given by the equation

$$z_{ij}^{m_i} = x_i y_{ij}$$

It suffices to prove that the normalization of $\times_j B_{ij}$ is nonsingular for all i . If $b \in D_i$ then since $\bigcup_j H_{ij} \cap D_i = \emptyset$, say $b \notin H_{i,0}$, and $y_{i,0}$ is a unit in $\mathcal{O}_{B,b}$. The normalization of the product $\times_j B_{ij}$ is locally defined by the equations

$$\begin{aligned} z_{i,1}^{m_i} &= x_i \\ \left(\frac{z_{i,j}}{z_{i,1}}\right)^{m_i} &= y_{i,j} \quad j = 2, \dots, \dim B. \end{aligned}$$

Since $x_i, y_{i,2}, \dots, y_{i,\dim B}$ are either units or local parameters in $\mathcal{O}_{B,b}$, it follows that the normalization of the product $\times_j B_{ij}$ is nonsingular at b . A similar situation happens when $b \notin D_i$, since then x_i is a unit in $\mathcal{O}_{B,b}$. Thus, B' is nonsingular and ramified with index m_i along D_i and H_{ij} . Replacing B by D_i or H_{ij} , we get that $p^*(\sum_i D_i + \sum_{i,j} H_{ij})_{\text{red}}$ is a divisor of normal crossings. $\square$

Let $\tilde{U}_B = B \setminus (\bigcup_{i,j} H_{ij} \cup D_i)$, and $\tilde{U}_{B'} = p^{-1}(\tilde{U}_B)$. Both $\tilde{U}_B \subset B$ and $\tilde{U}_{B'} \subset B'$ are toroidal embeddings because B, B' are nonsingular and the divisors $B \setminus \tilde{U}_B$, $B' \setminus \tilde{U}_{B'}$ cross normally. From the construction and local equations we also see that p is a toroidal morphism with respect to this structure. If $b' \in B'$, $b = p(b') \in B$, then there exist local parameters $\{x_l\}$ at b and $\{z_l\}$ at b' such that p is defined by $x_l = z_l^{a_l}$ where $a_l = m_i$ if the divisor defined by x_l is D_i or H_{ij} , otherwise $a_l = 1$. We denote the polyhedral complexes of $\tilde{U}_B \subset B$ and $\tilde{U}_{B'} \subset B'$ by $\tilde{\Delta}_B$ and $\tilde{\Delta}_{B'}$, respectively. So, every cone $\tau' \in \tilde{\Delta}_{B'}$ is mapped homeomorphically to a cone $\tau \in \tilde{\Delta}_B$ by p_Δ . If N_τ has basis $u_1, \dots, u_m$ then $N_{\tau'}$ has basis $a_1 u_1, \dots, a_m u_m$ where a_l are as above.

5.4. The toroidal picture. We need to see how adding the divisors H_{ij} to the toroidal structure of B affects the toroidal structure of X .

Lemma 5.7. *Let $f : X \rightarrow B$ be toroidal, B nonsingular, H a generic hyperplane section of B . Let $x \in X$, $b = f(x) \in H$. Then*

- (i) *there exist local models $(X_\sigma = X'_\sigma \times \mathbb{G}_m, z \times 1)$ at x and $(X_\tau = X'_\tau \times \mathbb{G}_m, y \times 1)$ at b such that H corresponds to the divisor $X'_\tau \times \{1\}$ and the morphism f is a product of toroidal morphisms*

$$f = g \times \text{id} : X'_\sigma \times \mathbb{G}_m \rightarrow X'_\tau \times \mathbb{G}_m$$

- (ii) *$U_B \setminus H \subset B$, $U_X \setminus f^{-1}(H) \subset X$ are toroidal embeddings and f is a toroidal morphism of these embeddings.*

Proof. Let (X_σ, s) and (X_τ, t) be local models at $x \in X$ and $b = f(x) \in B$, respectively, and let f also denote the toric morphism of the local models defined by

$$f_\Delta : (N_\sigma, \sigma) \rightarrow (N_\tau, \tau)$$

Clearly $U_B \setminus H \subset B$ is toroidal and we may assume that it has a local model $(X_{\langle \tau, v \rangle}, t')$, where $\langle \tau, v \rangle$ is the cone spanned by τ and some indivisible $v \in N_\tau$, and where t' lies in the unique closed orbit of $X_{\langle \tau, v \rangle}$. Write $N_{\tau,2}$ for the saturated sublattice of N_τ generated by v , and choose a splitting

$$\begin{aligned} N_\tau &\cong N_{\tau,1} \oplus N_{\tau,2} \\ X_\tau &\cong X'_\tau \times \mathbb{G}_m \\ X_{\langle \tau, v \rangle} &\cong X'_\tau \times \mathbb{A}^1 \end{aligned}$$

Since f is dominant, $\tilde{N}_{\tau,2} = f_\Delta(f_\Delta^{-1}(N_{\tau,2}))$ is a sublattice of finite index of $N_{\tau,2}$. The inclusion $\tilde{N}_{\tau,2} \subset N_{\tau,2}$ corresponds to an étale cover of $\mathbb{G}_m$. Since the completed local rings are isomorphic, we may replace the local model by a local model in the étale cover and assume that $f_\Delta^{-1}(N_{\tau,2})$ surjects onto $N_{\tau,2}$.

Let $N_{\sigma,1} = f_\Delta^{-1}(N_{\tau,1}) \subset N_\sigma$ and let $N_{\sigma,2} \subset N_\sigma$ be generated by some $u \in N_\sigma$ such that $f_\Delta(u) = v$. For any $w \in N_\sigma$, $f_\Delta(w) = v_1 + v_2 \in N_{\tau,1} \oplus N_{\tau,2}$, there exists $m \in \mathbb{Z}$ so that $f_\Delta(mu) = v_2$. Hence $w - mu \in N_{\sigma,1}$, and $N_\sigma = N_{\sigma,1} + N_{\sigma,2}$. Since $N_{\sigma,1} \cap N_{\sigma,2} = \{0\}$ the sum is direct,

$$\begin{aligned} N_\sigma &\cong N_{\sigma,1} \oplus N_{\sigma,2} \\ X_\sigma &\cong X'_\sigma \times \mathbb{G}_m \end{aligned}$$

and f_Δ maps $N_{\sigma,1}$ to $N_{\tau,1}$ and $N_{\sigma,2}$ to $N_{\tau,2}$. Thus the toric morphism f is a product $f = g \times h$ where $g : X'_\sigma \rightarrow X'_\tau$ and $h = id : \mathbb{G}_m \rightarrow \mathbb{G}_m$ are the toric morphisms induced by the restriction of f_Δ to $N_{\sigma,1}$ and $N_{\sigma,2}$, respectively. Since f_Δ maps $\langle \sigma, u \rangle$ to $\langle \tau, v \rangle$, f extends to a toric morphism

$$f : X_{\langle \sigma, u \rangle} \cong X'_\sigma \times \mathbb{A}^1 \xrightarrow{g \times id} X'_\tau \times \mathbb{A}^1 \cong X_{\langle \tau, v \rangle}$$

For $t = (y, 1) \in X'_\tau \times \mathbb{G}_m \subset X'_\tau \times \mathbb{A}^1$ the complete local rings $\hat{\mathcal{O}}_{X'_\tau \times \mathbb{G}_m, (y, 1)}$ and $\hat{\mathcal{O}}_{X'_\tau \times \mathbb{A}^1, (y, 0)}$ are isomorphic via translation. The closures of codimension 1 orbits through $(y, 0)$, are those through $(y, 1)$ plus $X'_\tau \times \{0\}$ which formally corresponds to H . Similarly, for $s = (z, 1) \in X'_\sigma \times \mathbb{G}_m \subset X'_\sigma \times \mathbb{A}^1$ the ring $\hat{\mathcal{O}}_{X'_\sigma \times \mathbb{G}_m, (z, 1)}$ is isomorphic to $\hat{\mathcal{O}}_{X'_\sigma \times \mathbb{A}^1, (z, 0)}$ via translation, and the closures of codimension 1 orbits through $(z, 0)$, are those through $(z, 1)$ plus $X'_\sigma \times \{0\} = f^*(X'_\tau \times \{0\})$ which formally corresponds to $f^*(H)$.

Thus $(X_{\langle \sigma, u \rangle}, (z, 0))$ and $(X_{\langle \tau, v \rangle}, (y, 0))$ are local models at $x \in X$ and $b \in B$, respectively, and the morphism

$$f : (U_X \setminus f^{-1}(H) \subset X) \rightarrow (U_B \setminus H \subset B)$$

is toroidal. □

It follows from the lemma that $f : (\tilde{U}_B \subset B) \rightarrow (\tilde{U}_X \subset X)$ is toroidal, where $\tilde{U}_X = f^{-1}(\tilde{U}_B)$. By Lemma 5.3, the normalization X' of $X \times_B B'$ is toroidal. Let $f' : X' \rightarrow B'$, $p' : X' \rightarrow X$ be the (toroidal) projections, and let $\hat{\Delta}_X, \hat{\Delta}_{X'}$ be the polyhedral complexes of X, X' .

$$\begin{array}{ccc} X' & \xrightarrow{p'} & X \\ f' \downarrow & & \downarrow f \\ B' & \xrightarrow{p} & B \end{array}$$

Then p'_Δ maps a cone $\sigma' \in \tilde{\Delta}_{X'}$ homeomorphically to a cone $\sigma \in \tilde{\Delta}_X$; the integral lattice of σ' can then be identified with a sublattice $N_{\sigma'} \subset N_\sigma$.

The following lemma shows that the added divisors $p^{-1}(H_{ij})$ and $(p \circ f')^{-1}(H_{ij})$ can be removed from the toroidal structures of B' and X' so that f' (but not p or p') remains toroidal.

Lemma 5.8. *$p^{-1}(U_B) \subset B'$ and $(p \circ f')^{-1}(U_B) \subset X'$ are toroidal embeddings and f' is a toroidal morphism of these embeddings.*

We show how to remove one irreducible divisor $H = H_{ij}$ for some i, j . Since the question is local, choose local models X_σ , X_τ and $X_{\tau'}$ of X , B , and B' so that both f and p are products

$$f : X_\sigma \cong X'_\sigma \times \mathbb{A}^1 \xrightarrow{g \times id} X'_\tau \times \mathbb{A}^1 \cong X_\tau$$

$$p : X_{\tau'} \cong X'_\rho \times \mathbb{A}^1 \xrightarrow{q \times r} X'_\tau \times \mathbb{A}^1 \cong X_\tau$$

where all morphisms are toric and H corresponds to $X'_\tau \times \{0\}$. Since p ramifies along H , the morphism $r : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ has degree $m \geq 2$.

The fiber product is then

$$\begin{aligned} X'' &= (X'_\sigma \times \mathbb{A}^1) \times_{X'_\tau \times \mathbb{A}^1} (X'_\tau \times \mathbb{A}^1) \\ &\cong (X'_\sigma \times_{X'_\tau} X'_\tau) \times (\mathbb{A}^1 \times_{\mathbb{A}^1} \mathbb{A}^1) \\ &\cong (X'_\sigma \times_{X'_\tau} X'_\tau) \times \mathbb{A}^1 \end{aligned}$$

and the projection $X'' \rightarrow X_{\tau'}$ is the product of the projection $X'_\sigma \times_{X'_\tau} X'_\tau \rightarrow X'_\tau$ and $id : \mathbb{A}^1 \rightarrow \mathbb{A}^1$. The divisor $(p \circ f')^*(H)$ in X'' is $X'_\sigma \times_{X'_\tau} X'_\tau \times \{0\}$ and by the same translation argument as above we can remove $p^*(H)$ and $(p \circ f')^*(H)$ from the toroidal structures of B' and X'' , respectively. $\square$

Let $\Delta_{X'}, \Delta_{B'}$ be the polyhedral complexes of the embeddings $p^{-1}(U_B) \subset B'$ and $(p \circ f')^{-1}(U_B) \subset X'$. Removing the divisors $p^{-1}(H_{ij})$ and $(p \circ f')^{-1}(H_{ij})$ means removing the corresponding edges (and everything attached to them) from the polyhedral complexes $\tilde{\Delta}_{B'}$ and $\tilde{\Delta}_{X'}$. As p_Δ and p'_Δ map cones of $\tilde{\Delta}_{B'}$ and $\tilde{\Delta}_{X'}$ homeomorphically to cones of $\tilde{\Delta}_B$ and $\tilde{\Delta}_X$, restrictions of p_Δ and p'_Δ map cones of $\Delta_{B'}$ and $\Delta_{X'}$ homeomorphically to cones of Δ_B and Δ_X . We summarize the previous constructions in the following proposition.

Proposition 5.9. *With the assumptions of Proposition 5.1, let u_i be the primitive points of Δ_B , and let $m_i > 0$ for $i = 1, \dots, l$. There exists a finite covering $p : B' \rightarrow B$ so that, if X' is the normalization of $X \times_B B'$ and $f' : X' \rightarrow B'$, $p' : X' \rightarrow X$ the two projections, we have*

- (i) $U_{X'} \subset X'$ and $U_{B'} \subset B'$ are toroidal embeddings with polyhedral complexes $\Delta_{X'}$ and $\Delta_{B'}$; moreover, f' is a toroidal morphism of these embeddings.
- (ii) There exist morphisms

$$p_\Delta : \Delta_{B'} \rightarrow \Delta_B$$

and

$$p'_\Delta : \Delta_{X'} \rightarrow \Delta_X,$$

such that

- for any cone $\tau' \in \Delta_{B'}$, the morphism p_Δ maps τ' isomorphically to τ , and identifies $N_{\tau'}$ with the sublattice of N_τ generated by $m_i u_i$;
- for any cone $\sigma' \in \Delta_{X'}$, p'_Δ maps σ' isomorphically to σ , and identifies $N_{\sigma'}$ with the sublattice $N_\sigma \cap f_\Delta^{-1}(N_{\tau'})$ of N_σ

◦ The following diagrams commute:

$$\begin{array}{ccc}
 \sigma' & \xrightarrow[p'_\Delta]{\cong} & \sigma \\
 f'_\Delta \downarrow & & \downarrow f_\Delta \\
 \tau' & \xrightarrow[p_\Delta]{\cong} & \tau
 \end{array}
 \qquad
 \begin{array}{ccc}
 N_{\sigma'} & \xrightarrow[p'_\Delta]{\subset} & N_\sigma \\
 f'_\Delta \downarrow & & \downarrow f_\Delta \\
 N_{\tau'} & \xrightarrow[p_\Delta]{\subset} & N_\tau
 \end{array}$$

□

Proof of 5.1 Let Δ_X, Δ_B be the polyhedral complexes of X, B , and $f_\Delta : \Delta_X \rightarrow \Delta_B$. By equidimensionality, f_Δ maps $\sigma^{(1)}$ to $\tau^{(1)}$. If u_i are the primitive points of Δ_B , let v_{ij} be the primitive points of Δ_X such that $f_\Delta(v_{ij}) = m_{ij}u_i$ for some $m_{ij} > 0$. Set $m_i = \text{lcm}_j\{m_{ij}\}$. Then for all i, j , some multiple of v_{ij} maps to $m_i u_i$. Now we use the covering data (D_i, m_i) to define the toroidal morphism $f' : X' \rightarrow B'$ as in the previous proposition. Let $\sigma' \in \Delta_{X'}$, $f'_\Delta(\sigma') = \tau' \in \Delta_{B'}$. There exist $\sigma \in \Delta_X, \tau \in \Delta_B$ such that $\sigma' \cong \sigma, \tau' \cong \tau$. The integral lattice $N_{\tau'}$ is generated by $m_i u_i$ with u_i lying on the edges of τ , and $N_{\sigma'} = N_\sigma \cap f_\Delta^{-1}(N_{\tau'})$. For any edge of σ' spanned by v_{ij} , some multiple $a_{ij}v_{ij}$ maps to $m_i u_i$. Hence $a_{ij}v_{ij} \in N_{\sigma'}$ is primitive and maps to the primitive point $m_i u_i$. This proves that f' has reduced fibers. □

6. MILDNESS OF THE MORPHISM

Recall that a normal variety X is said to have Gorenstein singularities if it is Cohen-Macaulay and has an invertible dualizing sheaf ω_X . It is said to have rational singularities, if for a resolution of singularities $r : X' \rightarrow X$ we have $r_*\omega_{X'} = \omega_X$. Every toric variety is Cohen-Macaulay with rational singularities ([KKMS] Theorem 14, p. 52). The dualizing sheaf of an affine toric variety X_τ is the coherent sheaf $\omega_{X_\tau} = \mathcal{O}(-\sum D_i)$ where D_i are closures of the codimension 1 orbits. This sheaf is invertible if and only if there exists an element of M (namely a linear function on τ taking integer values on N) such that $\psi(v) = -1$ for all $v \in \tau^{(1)}$.

Lemma 6.1. *Let X_σ and X_τ be affine toric varieties with X_τ nonsingular. Let $f : X_\sigma \rightarrow X_\tau$ be an equidimensional toric morphism without horizontal divisors, having only reduced fibers. Then X_σ has rational Gorenstein singularities.*

Proof. Since X_τ is nonsingular, it has rational Gorenstein singularities. Let $\psi : \tau \rightarrow \mathbb{R}$ be the linear interpolation of $\psi(u_i) = -1$ for every primitive point u_i of τ . Then $\psi \circ f_\Delta : \sigma \rightarrow \mathbb{R}$ is the required function. It is linear because ψ and f_Δ are; since f is equidimensional and has reduced fibers, primitive points in σ map to primitive points in τ , hence $\psi \circ f_\Delta$ takes the value -1 on every primitive point of σ . □

Lemma 6.2. *Let $U_X \subset X$ and $U_B \subset B$ be toroidal embeddings, and assume that B is nonsingular. Let $f : X \rightarrow B$ be an equidimensional toroidal morphism, without horizontal divisors, and with reduced fibers. Then X has rational Gorenstein singularities.*

Proof. Having rational Gorenstein singularities is a local analytic property. Since all local models have rational Gorenstein singularities, so does X . □

To show that the morphism is mild, we need to look at the situation after dominant base changes. We first look at the cases where the base change is relatively nice:

Lemma 6.3. *Let $f : X \rightarrow B$ be as above. Let $g : B' \rightarrow B$ be a dominant morphism, where B' a nonsingular variety, and assume that $g^{-1}(B \setminus U_B)$ is a normal crossings divisor. Let X' be the pullback of X to B' . Then $X' \rightarrow B'$ admits a toroidal structure relative to $g^{-1}U_B \subset B'$.*

Proof. We use the formalism of logarithmic structures. By [Kato1], §8.1, the morphism $X \rightarrow B$ is logarithmically smooth. Also $g^{-1}U_B \subset B'$ endows B' with a logarithmically regular structure. Moreover $g : B' \rightarrow B$ is a morphism of logarithmic schemes. The variety X' thus inherits a logarithmic structure. The morphism $X' \rightarrow B'$ clearly satisfies the formal lifting property for logarithmic smoothness ([Kato1], §8.1,(i)). It is left to show that X' satisfies condition (S) ([Kato1] §1.5). Indeed, since f is equidimensional, the fibers of f are reduced and B' is normal, it follows that X' is regular in codimension 1. Since f is a Gorenstein morphism and B is Gorenstein, we have that X' is Gorenstein, in particular it is Cohen Macaulay. It then follows that X' is normal. Combining this with the assumption that f^*M_B is saturated in M_X , we have that the monoids giving the logarithmic charts on X' are integral and saturated. Altogether, we have that X' satisfies condition (S). Therefore $f' : X' \rightarrow B'$ is logarithmically smooth, and thus toroidal.

Lemma 6.4. *Let $f : X \rightarrow B$ be a flat Gorenstein morphism and assume that B has rational Gorenstein singularities. Assume there is a modification $r : B' \rightarrow B$ with rational Gorenstein singularities, such that $X' = X \times_B B'$ has rational singularities. Then X has rational singularities as well.*

Proof. Consider the diagram

$$\begin{array}{ccc} X' & \xrightarrow{h} & X \\ f' \downarrow & & \downarrow f \\ B' & \xrightarrow{r} & B. \end{array}$$

By base change we have $h^*\omega_f = \omega_{f'}$. By assumption we have $\omega_B = r_*\omega_{B'}$. The flatness of f implies $f^*r_*\omega_{B'} = h_*f'^*\omega_{B'}$. Therefore we have

$$\begin{aligned} h_*\omega_{X'} &= h_*(\omega_{f'} \otimes f'^*\omega_{B'}) \\ &= h_*(h^*\omega_f \otimes f'^*\omega_{B'}) \\ &= \omega_f \otimes h_*f'^*\omega_{B'} \\ &= \omega_f \otimes f^*r_*\omega_{B'} \\ &= \omega_f \otimes f^*\omega_B \\ &= \omega_X. \end{aligned}$$

Thus X has rational singularities. □

Proposition 6.5. *Let $U_X \subset X$ and $U_B \subset B$ be toroidal embeddings and assume that B is nonsingular. Let $f : X \rightarrow B$ be an equidimensional morphism with reduced fibers, which is toroidal such that $U_X = f^{-1}U_B$. Then f is mild.*

Proof. Let $B_1 \rightarrow B$ be a dominant morphism such that B_1 has rational Gorenstein singularities. We need to show that $X_1 = X \times_B B_1$ has rational Gorenstein singularities. By lemma 6.2 it has Gorenstein singularities.

Pick a resolution of singularities $B' \rightarrow B_1$ such that the inverse image of $B \setminus U_B$ in B' is a divisor with normal crossings. By lemma 6.3 we have that $X' = X \times_B B'$ is toroidal, therefore X' has rational singularities. By lemma 6.4 we have that X_1 has rational singularities as well, which is what we needed. □

7. THE CARTIER COVERING

In section 8 below we will translate the semistable reduction conjecture in purely combinatorial terms. In order to maximize the flexibility of the combinatorial operations, we need to

generalize Kawamata's trick slightly to accommodate cases where B has quotient singularities. In such a situation we need a finite cover of B which is nonsingular, and such that the resulting polyhedral complex is easily described. The following statement will suffice for this purpose.

Proposition 7.1. *Let $U_X \subset X$ and $U_B \subset B$ be projective toroidal embeddings, and assume that B has only quotient singularities. Let $f : X \rightarrow B$ be a surjective mild toroidal morphism, satisfying $f^{-1}(U_B) = U_X$. Then there exists a finite surjective morphism $p : B' \rightarrow B$ so that, denoting $X' = X \times_B B'$, $U_{B'} = p^{-1}U_B$ and $U_{X'} = X \times_B U_{B'}$ we have*

1. B' and X' are toroidal embeddings with polyhedral complexes $\Delta_{X'}$ and $\Delta_{B'}$, with B' nonsingular;
2. the projection $f' : X' \rightarrow B'$ is a mild toroidal morphism;
3. There exist morphisms

$$p_\Delta : \Delta_{B'} \rightarrow \Delta_B$$

and

$$p'_\Delta : \Delta_{X'} \rightarrow \Delta_X,$$

such that

- for any cone $\tau' \in \Delta_{B'}$, the morphism p_Δ maps τ' isomorphically to τ , and identifies $N_{\tau'}$ with the sublattice of N_τ generated by the primitive vectors of τ ;
- for any cone $\sigma' \in \Delta_{X'}$, p'_Δ identifies $N_{\sigma'}$ with the sublattice $N_\sigma \cap f_\Delta^{-1}(N_{\tau'})$ of N_σ , and maps σ' isomorphically to σ .
- The following diagrams commute:

$$\begin{array}{ccc} \sigma' & \xrightarrow[p'_\Delta]{\cong} & \sigma \\ f'_\Delta \downarrow & & \downarrow f_\Delta \\ \tau' & \xrightarrow[p_\Delta]{\cong} & \tau \end{array} \qquad \begin{array}{ccc} N_{\sigma'} & \xrightarrow[p'_\Delta]{\subseteq} & N_\sigma \\ f'_\Delta \downarrow & & \downarrow f_\Delta \\ N_{\tau'} & \xrightarrow[p_\Delta]{\subseteq} & N_\tau \end{array}$$

Proof. We use a construction analogous to Kawamata's covering. First, let B be a variety, E an effective Weil divisor on B . We make the following assumptions:

1. as a scheme, E is integral and normal;
2. the divisor $mE = D$ is a Cartier;
3. There is a Cartier divisor L on B such that $\mathcal{O}(D) = \mathcal{O}(mL)$.

As in Definition 5.4, we choose a function ϕ with $\text{div}(\phi) = D - mL$, and define the cyclic cover $p : B' \rightarrow B$ by taking m -th root of ϕ . The point is, that the $\mathbb{Q}$ -Cartier divisor p^*E is in fact Cartier.

Let us see what happens in the toric case.

Lemma 7.2. *Let X_τ be an affine toric variety with quotient singularities and E a toric divisor corresponding to a primitive vector v . Write $\tau = \mathbb{R}_+ v \times \tau_1$. Choose $m \in \mathbb{N}$ such that $D = mE$ is Cartier. Then $\mathcal{O}(mD)$ is trivial, and the corresponding cyclic cover is $X_{\tau'}$, where $|\tau'| = \tau$ and $N_{\tau'} = \mathbb{Z}v \times (N_\tau \cap \tau_1)$.*

This lemma is easy and left to the reader. If we iterate the lemma with all the toric divisors, we obtain a nonsingular covering, whose lattice is generated by the primitive vectors of τ .

Assume that in addition we have a mild morphism $X_\sigma \rightarrow X_\tau$. Lemma 5.3 works word-for-word in this case. The only thing which needs to be changed in the proof is, that since $f_\Delta(N_\sigma) = N_\tau$, the semigroup homomorphism $i : \pi \rightarrow M$ is still injective. (This is not a serious business - we could replace the product by its reduction anyway.)

In order to define a Kawamata covering package, we need the following Bertini type lemma (this is a special case of the stratified Bertini Theorem) .

Lemma 7.3. *Let $U_B \subset B$ be a toroidal embedding. Let H_0 be a very ample divisor on B and let H be a general element of $|H_0|$. Let $U'_B = U_B \setminus H$. We have*

1. $U'_B \subset B$ is a toroidal embedding;
2. let $b \in H$ and assume that b lies on a stratum $Y \subset B$ of the toroidal embedding $U_B \subset B$. Then $H \cup Y$ is nonsingular at b ;
3. there is a local model $X_\rho \times \mathbb{G}_m^k$ for $U_B \subset B$ at b , where Y corresponds to the factor $\mathbb{G}_m^k$, and H corresponds to $X_\rho \times \mathbb{G}_m^{k-1} \times \{1\}$.
4. there is a local model $X_\rho \times \mathbb{G}_m^{k-1} \times \mathbb{A}^1$ for $U'_B \subset B$ at b , where Y corresponds to $\mathbb{G}_m^k \times \mathbb{A}^1$, and H corresponds to $X_\rho \times \mathbb{G}_m^{k-1} \times \{0\}$.

Proof. Since B is a stratified space, it has a local product structure $X_\rho \times Y$ at b . We need to adapt this structure to the divisor H as in the lemma. Applying Bertini's theorem to the closure of the stratum Y , we have that $H \cup Y$ is nonsingular. Pick a regular function u near b defining H on a neighborhood of b . Then the restriction u_Y of u to Y is a regular parameter. We can find a regular system of parameters on Y at b including u_Y and lift them to B . This lifting gives the desired adaptation of the product structure. $\square$

Using this lemma, the proof of Proposition 5.9 works word-for-word here, and yields Proposition 7.1. $\square$

8. TOWARDS SEMISTABLE REDUCTION

8.1. Combinatorial semistable reduction. We are going to restate the semistable reduction conjecture (Conjecture 0.2) in purely combinatorial terms. First let us define semistability combinatorially.

Definition 8.1. Let Δ_X and Δ_B be rational conical polyhedral complexes, and let $f_\Delta : \Delta_X \rightarrow \Delta_B$ be a surjective polyhedral map. We say that f_Δ is **weakly semistable** if the following conditions hold.

1. We have $f_\Delta^{-1}(0) = \{0\}$.
2. For any cone $\tau \in \Delta_X$, we have $f_\Delta(\tau) \in \Delta_B$.
3. For any cone $\tau \in \Delta_X$, we have $f_\Delta(N_\tau) = N_{f_\Delta(\tau)}$.
4. Δ_B is simplicial with index 1.

If also Δ_X is simplicial with index 1, we say that f_Δ is **semistable**.

We now define the operations we are allowed to perform.

Definition 8.2. Let Δ be a rational conical polyhedral complex. A **lattice alteration** $\Delta_1 \rightarrow \Delta$ consists of an integral structure induced by a consistent choice of sublattices $N_\sigma^1 \subset N_\sigma$ for each cone $\sigma \subset \Delta$. An **alteration** $\Delta_1 \rightarrow \Delta$ is a composition $\Delta_1 \rightarrow \Delta' \rightarrow \Delta$ of a lattice alteration $\Delta_1 \rightarrow \Delta'$ with a subdivision $\Delta' \rightarrow \Delta$. The alteration is **projective** if the corresponding subdivision $\Delta' \rightarrow \Delta$ is projective.

Note that the composition of alterations is an alteration. Be warned that the factorization of a polyhedral alteration above is not analogous to the Stein factorization in algebraic geometry - it has the opposite order.

Definition 8.3. 1. Let $\Delta_X \rightarrow \Delta_B$ be a polyhedral map of rational conical polyhedral complexes. Let $\Delta'_B \rightarrow \Delta_B$ be a subdivision. The induced subdivision $\Delta'_X \rightarrow \Delta_X$ is the

- minimal subdivision admitting a polyhedral map to Δ'_B . The cones of Δ'_X are of the form $\sigma \cap f_\Delta^{-1}(\tau')$ for $\sigma \in \Delta_X$ and $\tau' \in \Delta'_B$.
2. Let $\Delta_X \rightarrow \Delta_B$ be a polyhedral map of rational conical polyhedral complexes. Let $\Delta_B^1 \rightarrow \Delta_B$ be a lattice alteration. The induced lattice alteration $\Delta_X^1 \rightarrow \Delta_X$ is the minimal lattice alteration mapping to Δ_B^1 . The lattices of Δ_X^1 are of the form $N_\sigma \cap f_\Delta^{-1}(N_{\tau^1})$ for $\sigma \in \Delta_X$ and $\tau^1 \in \Delta_B^1$.
 3. Let $\Delta_X \rightarrow \Delta_B$ be a polyhedral map of rational conical polyhedral complexes. Let $\Delta_B^1 \rightarrow \Delta'_B \rightarrow \Delta_B$ be an alteration, factored as a lattice alteration followed by a subdivision. The induced alteration $\Delta_X^1 \rightarrow \Delta_X$ is the induced lattice alteration $\Delta_X^1 \rightarrow \Delta'_X$ of the induced subdivision $\Delta'_X \rightarrow \Delta_X$.

Note that an alteration induced by a projective alteration is projective.

We are now ready to state our conjecture.

Conjecture 8.4. Let $f_\Delta : \Delta_X \rightarrow \Delta_B$ be a polyhedral map of rational conical polyhedral complexes, and assume for simplicity that $f_\Delta^{-1}(0) = \{0\}$. Then there exists a projective alteration $\Delta_B^1 \rightarrow \Delta_B$, with induced alteration $\Delta_X^1 \rightarrow \Delta_X$, and a projective subdivision $\Delta_Y \rightarrow \Delta_X^1$, such that $\Delta_Y \rightarrow \Delta_B^1$ is semistable.

The conjectures are tied together by the following proposition:

Proposition 8.5. *Conjecture 8.4 implies Conjecture 0.2.*

Proof. Let $X \rightarrow B$ be a surjective morphism of complex projective varieties with geometrically integral generic fiber. By Theorem 2.1 we may assume the morphism toroidal. Let $f_\Delta : \Delta_X \rightarrow \Delta_B$ be the associated polyhedral map. By theorem 0.3 we may assume f_Δ is weakly semistable. We now invoke Conjecture 8.4. Let $\Delta_B^1 \rightarrow \Delta'_B \rightarrow \Delta_B$ be a projective alteration, $\Delta_X^1 \rightarrow \Delta_X$ the induced alteration, and $\Delta_Y \rightarrow \Delta_X^1$ a projective subdivision such that $\Delta_Y \rightarrow \Delta_B^1$ is semistable. The subdivision $\Delta'_B \rightarrow \Delta_B$ and the induced subdivision $\Delta'_X \rightarrow \Delta_X$ give rise to modifications $B' \rightarrow B$ and $X' \rightarrow X$ with a lifting $X' \rightarrow B'$ (see lemma 1.7). Let $\Delta_B^0 \rightarrow \Delta'_B$ be the lattice alteration given by the sublattice generated by the primitive vectors, and $\Delta_X^0 \rightarrow \Delta'_X$ the induced lattice alteration. Proposition 7.1 gives rise to an alteration $\widetilde{B}_0 \rightarrow B'$, with pullback $\widetilde{X}_0 \rightarrow X'$, and a polyhedral map $\Delta_{\widetilde{B}_0} \rightarrow \Delta_B^0$ which is an isomorphism on each cone. The same holds for $\Delta_{\widetilde{X}_0} \rightarrow \Delta_X^0$. The lattice alteration $\Delta_B^1 \rightarrow \Delta'_B$, being of index 1, factors through Δ_B^0 . Proposition 5.9 gives an alteration $\widetilde{B}_1 \rightarrow \widetilde{B}_0$ with pullback $\widetilde{X}_1 \rightarrow \widetilde{X}_0$ and a polyhedral map $\Delta_{\widetilde{B}_1} \rightarrow \Delta_{\widetilde{B}_0}^1$ which is an isomorphism on each cone, and the same holds for $\Delta_{\widetilde{X}_1} \rightarrow \Delta_{\widetilde{X}_0}^1$. The subdivision $\Delta_Y \rightarrow \Delta_X^1$ induces a subdivision $\widetilde{\Delta}_Y \rightarrow \Delta_{\widetilde{X}_1}^1$, giving rise to a toroidal modification $\widetilde{Y} \rightarrow \widetilde{X}_1$. Since $\Delta_Y \rightarrow \Delta_B^1$ is semistable, we have that $\widetilde{\Delta}_Y \rightarrow \Delta_{\widetilde{B}_1}^1$ is semistable as well, and therefore $\widetilde{Y} \rightarrow \widetilde{B}_1$ is semistable, as required. $\square$

8.2. How far can we push the results? In [N-R], it is shown that if $\Delta_0 \subset \Delta$ is a subcomplex, and $\Delta'_0 \rightarrow \Delta_0$ is a *projective* triangulation, then there is a projective triangulation $\Delta' \rightarrow \Delta$ extending Δ'_0 , without new edges: $\Delta'^{(1)} = \Delta^{(1)} \cup \Delta'^{(1)}$. Applying the main theorem of [KKMS], Chapter III, it is shown in [N-R] that given a weakly semistable $X \rightarrow B$, one can locally find a finite map $B_1 \rightarrow B$ and a modification $X_2 \rightarrow X \times_B B_1$ such that $X_2 \rightarrow B_1$ is semistable in codimension 1. If we apply Kawamata's trick, then clearly this can be done globally. Moreover, X_2 has only quotient singularities.

A weakly semistable morphism is said to be **almost semistable** if it is semistable in codimension 1 and moreover X has quotient singularities. Thus, in theorem 0.3 we may replace

“weakly semistable” by “almost semistable”. An analogous definition can be made on the polyhedral side.

It is important to note that an almost semistable morphism is not necessarily semistable. It is easy to give a polyhedral example: let $\tau = (\mathbb{R}_+)^2$ be endowed with the standard lattice $\mathbb{Z}^2$, and let $\sigma = (\mathbb{R}_+)^4$ be given the lattice generated by $\mathbb{Z}^4$ and the vector $w = (1/2, 1/2, 1/2, 1/2)$. We have a polyhedral map $\sigma \rightarrow \tau$ given by $(a, b, c, d) \mapsto (a + b, c + d)$. It is easy to see that this is almost semistable, but not semistable, since σ has index 2. Needless to say, a corresponding toroidal example can be easily constructed as well.

The example we just gave is easy to amend. Indeed, if we subdivide τ at its barycenter $(1, 1)$, take the induced subdivision of σ , then its star subdivision centered at w , and extend this to a triangulation using [N-R], we obtain a semistable map. This can be extended to families of surfaces in general - the main observation (see [Wang] for the case $\dim(B) = 1$) is that one can use Pick’s theorem and subdivide, with no need for additional lattice alteration. We plan to pursue this elsewhere.

One last remark: the second author has shown, that in order to prove semistable reduction, it is sufficient to produce B_1 and Y such that $Y \rightarrow B_1$ satisfies all but condition 3 of the requirements for semistability in Definition 8.1. Again, this will be pursued elsewhere.

REFERENCES

- [N] D. Abramovich, *A high fibered power of a family of varieties of general type dominates a variety of general type*, Inventiones Math., to appear.
alg-geom/9604024
- [N-dJ] D. Abramovich and A. J. de Jong, *Smoothness, semistability and toroidal geometry*, J. Alg. Geom., to appear.
alg-geom/9603018
- [N-R] D. Abramovich and J. M. Rojas, *Extending triangulations and semistable reduction*, preprint.
- [Al] V. Alexeev, *Boundedness and K^2 for log surfaces*. Internat. J. Math. 5 (1994), no. 6, 779-810.
- [Al1] V. Alexeev, *Moduli spaces $M_{g,n}(W)$ for surfaces*, preprint.
alg-geom/9410003
- [A-W] M. Artin and G. Winters, *Degenerate fibers and reduction of curves*, Topology 10 (1971) p. 373–383.
- [B-M] E. Bierstone and P. Milman, *Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant*, Invent. Math. 128 (1997), no. 2, 207–302.
- [D] V. I. Danilov, *The Geometry of Toric Varieties*, Russian Math. Survey, 33:2 (1978), 97-154.
- [dJ] A. J. de Jong, *Smoothness, semistability, and alterations*, Publ. Math. I.H.E.S. **83**, 1996, pp. 51-93.
- [dJ2] A. J. de Jong, *Families of curves and alterations*, Annales de l’Institut Fourier, to appear.
- [Kawa] Y. Kawamata, *Characterization of Abelian Varieties*, Compositio mathematica, vol. 43, Fasc. 2, 253-276 (1981).
- [Kato] K. Kato, *logarithmic structures of Fontaine-Illusie*, Algebraic analysis, geometry and number theory: Proceedings of the JAMI Inaugural Conference, 191–224, Johns Hopkins Univ. Press, Baltimore, MD, 1989.
- [Kato1] K. Kato, *Toric singularities*. Amer. J. Math. 116 (1994), no. 5, 1073–1099.
- [KKMS] G. Kempf, F. Knudsen, D. Mumford and B. Saint-Donat, *Toroidal Embeddings I*, Springer, LNM 339, 1973.
- [Mor] R. Morelli, *The birational geometry of toric varieties*. J. Alg. Geom. **5** (1996) 751-782.
- [Vie] E. Viehweg, *Quasi-projective moduli of polarized manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete Band 30. Springer, 1995.
- [Vil] O. Villamayor, *Constructiveness of Hironaka’s resolution*. Ann. Sci. École Norm. Sup. (4) 22 (1989), no. 1, 1–32.
- [Wang] J.-H. Wang, MIT dissertation.
- [Włó] J. Włodarczyk, *Decomposition of birational toric maps in blow-ups and blow-downs*. Trans. Amer. Math. Soc. 349 (1997), no. 1, 373–411.

DEPARTMENT OF MATHEMATICS, BOSTON UNIVERSITY, 111 CUMMINGTON, BOSTON, MA 02215, USA

E-mail address: `abrmovic@math.bu.edu`

E-mail address: `kllkr@math.bu.edu`